

METHODS AND COMPOSITIONS FOR TREATING  
ADDICTION DISORDERS

5

ABSTRACT OF THE DISCLOSURE

10051770-011802

A method for treating an addiction disorder (such as an addiction to or dependency on stimulants, nicotine, morphine, heroin, other opiates, amphetamines, cocaine, and/or alcohol) in a patient is disclosed. The method includes administering to the patient a first  $\alpha_3\beta_4$  nicotinic receptor antagonist and administering to the patient a second  $\alpha_3\beta_4$  nicotinic receptor antagonist. The second  $\alpha_3\beta_4$  nicotinic receptor antagonist is different than the first  $\alpha_3\beta_4$  nicotinic receptor antagonist, and the first  $\alpha_3\beta_4$  nicotinic receptor antagonist and the second  $\alpha_3\beta_4$  nicotinic receptor antagonist are administered simultaneously or non-simultaneously. Compositions which include a first  $\alpha_3\beta_4$  nicotinic receptor antagonist and a second  $\alpha_3\beta_4$  nicotinic receptor antagonist are also described. Examples of suitable  $\alpha_3\beta_4$  nicotinic receptor antagonists for use in the present invention's methods and compositions include mecamlamine, 18-methoxycoronaridine, bupropion, dextromethorphan, dextrorphan, and pharmaceutically acceptable salts and solvates thereof. A method of evaluating a compound for its effectiveness in treating addiction disorders is also described.